

Original Research

A retrospective study of road traffic accidents - injury characteristics, management and outcome at Rohilkhand Medical College and Hospital, Bareilly

Seth Sharad, Agarwal Keshav, Agarwal Parnnika

Department of Surgery, Rohilkhand Medical College and Hospital, Bareilly, Uttar Pradesh, India

ABSTRACT

Aim: This is a retrospective study carried out in Rohilkhand Medical College and Hospital, Bareilly to study the injury characteristics, management and outcome of road traffic accidents and factors responsible for these injuries.

Background: Urbanization has led to an increased number of vehicles on the roads. This together with an extreme apathy on the part of authorities to implement traffic rules has led to a phenomenal rise in the number of road traffic accidents. Lack of a proper pre-hospital care system has only added to the woes. Thus, the need for this study.

Materials and Methods: All case records of road traffic accident victims admitted to the hospital between January and December 2013 were analyzed statistically with respect to their demographic profile, injury characteristics, management and outcome. The revised trauma score was used to assess the patients and study its reliability in predicting mortality at the time of presentation to the casualty. **Results:** This study showed that the majority of crash victims were males in the age group 21-40 years accounting for huge economic losses for their families and the country at large. Motorized two-wheeler accounted for 62.5% of the accidents. Musculoskeletal injuries (57.5%) were the most common injuries, followed by those to the face, neck, and head. Local wound treatment (72%) was the most common surgical procedure performed. About 12.5% of the trauma victims did not survive their injury. **Conclusion:** Road traffic accidents are preventable. Strict traffic laws and penalties have to be imposed to curb this ever growing menace.

Keywords: Accidents, Wounds and Injuries, Treatment outcomes

INTRODUCTION

Road traffic accidents are a human tragedy. Nearly 1.3 million people die every year on the world's roads and up to 50 million people suffer non-fatal injuries, with many sustaining a disability as a result of their injury.¹ These injuries and deaths have an immeasurable impact on the families affected.

The United Nations has declared 2011-20 as the decade of action on road safety so that the present rising trend of road accidents stabilizes and is reversed by the year 2020.²

In India, road traffic accidents accounted for 2% of all deaths in 2005 nationally, of which 65% occurred in men between the ages of 15 and 59 years taking a heavy toll on those in their most productive years.³ 82.2% of road casualties were in the age group 15-65 years in the year 2011. Only 15% of the road accident victims were females in the calendar year 2012.¹

Road traffic accidents are both predictable and preventable. In India, there is no central authority to lay down guidelines for trauma care. There are no legislations in place to regulate qualification of ambulance personnel and maintenance of ambulance equipment. Rural India lacks appropriate trauma care facilities due to financial constraints and an underdeveloped health infrastructure. The government, medical fraternity and the society at large are yet to recognize trauma as a growing health hazard.⁴

Only a good combination between a mature pre-hospital system and a trauma care center can provide professional care and the best possible outcomes in cases of trauma.⁵

Lack of local data and information on road traffic accidents in our region has resulted in an utter disregard to this problem amongst our policy makers.

Corresponding Author:

Dr. Sharad Seth, Department of Surgery, Rohilkhand Medical College and Hospital, Bareilly, Uttar Pradesh, India.

E-mail: sharad_seth2002@yahoo.co.in

© 2015 International Journal of Medical Science Research and Practice available on www.ijmsrp.com

This study was conducted in order to identify the injury characteristics of road traffic accident victims and to provide baseline data to authorities who may wish to undertake interventions to improve road safety in the area.

MATERIALS AND METHODS

Place of Study

The study was carried out in The Department of Surgery, Rohilkhand Medical College and Hospital, Bareilly, during a period of 2 months from May 1st, 2014 to June 30th, 2014.

Study Design

This is a retrospective study of all road traffic accident patients, who presented in the casualty of Rohilkhand Medical College and Hospital, Bareilly, during the period January 2013-December 2013. Only those cases of road traffic accidents, which took place on the road and which involved any kind of moving vehicle and who reported directly to the casualty department of the hospital were considered.

Ethical Clearance

Approval from the Institutional Ethics Committee was obtained before commencement of the study.

Sample Size

Data of 120 patients were analyzed in this study.

Methodology of Data Collection

All case records were analyzed with respect to the age and sex of the patient, mechanism of injury, characteristics of the injury, management and outcome. The revised trauma score (RTS) was used to score and assess the condition of the patient, to see whether it had any reliability in predicting death. The score uses three variables namely Glasgow coma scale (GCS), systolic blood pressure (SBP), and respiratory rate (RR). All three indices are calculated based on a coded value. The score has values ranging from 0 to 7.8408.

GCS	SBP	RR	Coded value
13-15	>89	10-29	4
9-12	76-89	>29	3
6-8	50-75	6-9	2
4-5	1-49	1-5	1
3	0	0	0

GSC: Glasgow coma scale, SBP: Systolic blood pressure, RR: Respiratory rate

$$RTS = 0.9368 \text{ GCS} + 0.7326 \text{ SBP} + 0.2908 \text{ RR}$$

Data Analysis

All variables from the case study sheet were analyzed statistically. To study the relationship between RTS and mortality of patients, Chi-square test was used.

RESULTS

The maximum number of victims (22.5%) of RTA was in the age group of 21-40 years coupled with a male

preponderance. Most accidents (47.5%) occurred between 16:01 and 00:00 h, followed by 42.5% in the time interval between 08:01 and 16:00 h. In 63 cases (52.5%), the victim injured was the driver of the colliding vehicle.

Most accidents occurred in the city (45%), followed by 37.5% in the village. It was further seen that out of 120 victims, 105 (87.5%) were brought to the hospital by self/relatives/friends, 10 (8.34%) by good Samaritans and 5 (4.16%) by a government agency. Musculoskeletal injuries were the most common (57.5%). The face and neck accounted for (50%) of the injuries. 75% cases had superficial injuries like contusion/abrasion and 57.5% cases had lacerations. Fracture/dislocation was seen in 40% cases (Figure 1). Local wound treatment was required singly or additionally in 60% cases. The RTS was calculated for the victims retrospectively, it was seen that 33 victims had a score between 4.1 and 6, out of whom 21 patients survived and 12 died. 87 patients had a score >6, of whom, 84 patients survived and 3 died (Table 1). The Chi-square value came out to be 23.69 ($P < 0.001$).

It was seen that 90% of patients (108) required hospital admission. 10% patients (12) could be managed on an outpatient basis. Out of the total number of victims, 63 (52.5%) were discharged with no residual disability, 42 (35%) sustained some form of disability and 15 (12.5%) patients died.

DISCUSSION

A total of 120 cases of road traffic accidents were studied. Majority of victims (45%) were in the age group 21-40 years with a male preponderance. This is in concordance with other studies.^{6,7,8} Thus, accidents impacted the most productive age group with devastating consequences for the

Table 1: RTS of victims

RTS	Number of victims	Alive	Dead
4.1-6	33	21	12
>6	87	84	3

RTS: Revised trauma score




Figure 1: Injury characteristics among victims* (*multiple sites and injuries to patients were grouped in individual categories)

family. Reckless driving and speeding in this age group is responsible for most of the accidents.

Lack of adequate street lighting, bad conditions of roads, obstacles, unmarked speed-breakers, stray animals, no lights on bicycles and bullock-carts were probably responsible for most of the accidents occurring between 4:01 pm and 00:00 am (47.5%) as seen in another study.⁹ A significant number of accidents (42.5%) did occur in the day hours, 8 am to 4 pm. A hurry to get to work with scant regard for traffic regulations accounted for most of these cases.

Motorized two-wheeler were seen to be involved in the majority of collisions (62.5%), which is similar to those found in other studies.^{6,7,10} Increasing urbanization, easy availability of loans, affordability and catchy television advertisements has resulted in an ever increasing number of these vehicles on roads, which by their inherent instability and overloading make them extremely accident prone.

In the present study, it was found that the majority of victims were drivers of the vehicle involved in collision. This is in concordance with the study carried out in Lucknow by Chauhan *et al.*⁸ but it is in contrast to a study carried out in Tanzania by Chalya *et al.*,⁷ where pedestrians constituted the majority of victims. In a study by Jha *et al.* in Puducherry,¹¹ occupants of vehicles were mainly involved.

Our study shows that the most number of accidents occurred within the city (45%), followed by (37.5%) in the village. Chauhan *et al.* showed a similar result in their study carried out in Lucknow whereas Singh *et al.*, in their study in Ghaziabad region reported that maximum accidents occurred on the highway.^{7,12} The location of our hospital well within the city and best equipped to deal with accidents may be the reason for the above.

It was seen in this study that majority of victims (87.5%) came to the hospital by themselves or were brought by relatives/friends, 8.33% were brought by good Samaritans and only 4.16% by a government agency. Similar results were obtained by Chalya *et al.* in their study.⁷ The existing systems for pre-hospital care of trauma patients is still in its infancy in most parts in the country. Apathy toward the trauma victim coupled with complex legal issues deters people coming forward to help trauma victims. Ambulance drivers and other helpers have no formal training in trauma care. Even though the state government runs a free ambulance service, knowledge of its existence is not universally present among the village folk who still use local transport to shift patients.

In our study, musculoskeletal injuries to the extremities were the most common (57.5%), followed by face and neck (50%) and head (42.5%). Bhuyan and Ahmed in their study in Assam showed a greater number of head and neck injuries along with extremity injuries.¹³ Chalya *et al.* and Chauhan *et al.* in their study found a greater number of musculoskeletal and head injuries but a smaller number of maxillofacial trauma.^{7,8} The increased number of musculoskeletal injuries is obviously due to a fall from a two-wheeler. Even though, legislation for the compulsory usage of crash helmets exists, the public at large in this part of the

country disregards it accounting for the high numbers of head and maxillofacial injuries.

It was seen that superficial injuries like contusion and abrasion occurred in 75% and lacerations occurred in 57.5% of victims. Similar results were seen in a study carried out in Ghaziabad.¹² This is due to the obvious effect of blunt trauma and fall on the roads with a rough surface. Fracture/dislocation occurred in 40% of victims. Fractures were the most common injuries in studies carried out elsewhere.^{6,8} There is general lack of awareness among people regarding first aid treatment. This leads to increased manipulation of the injured limb worsening the nature of fracture. Local wound treatment was the most common surgical procedure performed in our study, followed by treatment of fracture/amputation. This is in concordance with the study carried out in Tanzania.⁷

In our study, the simple RTS was calculated for each victim retrospectively to study whether it would be useful in predicting death. It was seen that out of the 33 victims with score between 4.1 and 6, 21 patients survived and 12 died and, of the 87 patients with score >6, 84 patients survived and 3 died. The chi square value was 23.69 ($P < 0.001$). Thus, there was significant relationship between the RTS of victims and their final outcome, a lower score implying a poor prognosis. Other studies have also found the RTS as a simple and reliable indicator of prognosis in a trauma patient.¹⁴

In our study, 90% of the patients were managed after admission and only 10% of patients were managed on an outpatient basis. This is probably because local practitioners deal with most of the patients with minor injuries.

CONCLUSION

Road traffic accidents are an emerging problem in our nation. The most commonly affected are males in a young age group, which means that lives of many families are irrevocably changed by these mishaps. Motorized two wheelers are the offending vehicles in most cases.

Strict laws regarding speed limit, use of safety belts and helmets together with better quality roads and roadside illumination will go a long way in preventing such tragedies. A multi-disciplinary approach consisting of public education, a proper pre-hospital trauma care system and definitive trauma care facilities coupled with rehabilitation is required to be put in place if any impact is envisaged on this ever-growing epidemic on the roads.

ACKNOWLEDGMENTS

We are extremely grateful for the support received from the ICMR under their STS program in completion of this research.

PEER REVIEW

None.

CONFLICTS OF INTEREST

The authors declare that they have no competing interests.

FUNDING

Not applicable

REFERENCES

1. Ruikar M. National statistics of road traffic accidents in India. *J Orthop Traumatol Rehabil* 2013;6:1-6.
2. United Nations Decade of action for road safety 2011-2020. Available from: <http://www.decadeofaction.org>. [Last accessed on 2015 Mar 15].
3. Hsiao M, Malhotra A, Thakur JS, Sheth JK, Nathens AB, Dhingra N, *et al.* Road traffic injury mortality and its mechanisms in India: Nationally representative mortality survey of 1.1 million homes. *BMJ Open* 2013;3:e002621.
4. Joshipura MK, Shah HS, Patel PR, Divatia PA. Trauma care systems in India: An overview. *Indian J Crit Care Med* 2004;8:93-7.
5. Beuran M, Paun S, Gaspar B, Vartic N, Hostiuc S, Chiotoroiu A, *et al.* Prehospital trauma care: A clinical review. *Chirurgia (Bucur)* 2012;107:564-70.
6. Ganveer GB, Tiwari RR. Injury pattern among non-fatal road traffic accident cases: A cross-sectional study in Central India. *Indian J Med Sci* 2005;59:9-12.
7. Chalya PL, Mabula JB, Dass RM, Mbelenge N, Ngayomela IH, Chandika AB, *et al.* Injury characteristics and outcome of road traffic crash victims at Bugando Medical Centre in Northwestern Tanzania. *J Trauma Manag Outcomes* 2012;6:1.
8. Chauhan A, Ahmed N, Singh JV, Singh VK, Singh A, Kumar S. Epidemiology of road traffic injuries in a tertiary care Centre of Lucknow. *Indian J Community Health* 2014;26:181-6.
9. Dsouza C, Rao VV, Kumar A, Diaz E. Epidemiological trends of trauma in tertiary care Centre in Dakshina Kannada district of Karnataka, India. *J Clin Diagn Res* 2014;8:66-8.
10. Chandra Shekar BR, Reddy C. A five-year retrospective statistical analysis of maxillofacial injuries in patients admitted and treated at two hospitals of Mysore city. *Indian J Dent Res* 2008;19:304-8.
11. Jha N, Srinivasa DK, Roy G, Jagdish S. Injury pattern among road traffic accident cases: A study from South India. *Indian J Community Med* 2003;28:85.
12. Singh R, Bhatnager M, Singh HK, Singh GP, Kumar Y. An epidemiological study of victims of road traffic accidents cases: A study from national capital region (Ghaziabad), India. *Indian J Prev Soc Med* 2011;42:30-4.
13. Bhuyan PJ, Ahmed F. Road traffic accident: An emerging public health problem in Assam. *Indian J Community Med* 2013;38:100-4.
14. Champion HR, Sacco WJ, Copes WS, Gann DS, Gennarelli TA, Flanagan ME. A revision of the Trauma Score. *J Trauma* 1989;29:623-9.

How to cite this article: Sharad S, Keshav A, Parnnika A. A retrospective study of road traffic accidents - injury characteristics, management and outcome at Rohilkhand Medical College & Hospital, Bareilly. *Inter J Medical Sci Res Prac* 2015;2(2):73-76.

Received: 23 Mar 2015; **Accepted:** 30 Apr 2015; **Published:** 30 Jun 2015