

Image

Cutis marmorata telangiectatica congenita in an infant

Priya Gogia, Ravi Ambey

Department of Pediatrics, Gajra Raja Medical College, Gwalior, Madhya Pradesh, India



Figure 1: Infant with cutis marmorata telangiectatica congenita having marbled, reticulated vascular pattern of lower limb and trunk with a sharp cut-off line at midline

A 3-month-old male infant was brought to the hospital with the complaint of the reticulated pattern over the skin of limb and trunk. The pattern was present since birth. The blue - purplish marbled, reticulated pattern of the skin was present on the right lower limb and trunk with a sharp cutoff line at midline (Figure 1). The pattern was having multiple indentations, which is the characteristic feature of cutis marmorata telangiectatica congenita (Figure 2). The



Figure 2: Infant with cutis marmorata telangiectatica congenita showing characteristics indentation of skin

pattern was getting prominent on crying and cold exposure. Ophthalmic examination was normal. There was no associated systemic abnormality.

How to cite this article: Gogia P, Ambey R. Cutis marmorata telangiectatica congenita in an infant. Inter J Medical Sci Res Prac 2015;2(2):112-112.

Received: 06 Apr 2015; **Accepted:** 14 Jun 2015; **Published:** 30 Jun 2015

Corresponding Author:

Dr. Ravi Ambey, Department of Pediatrics, Gajra Raja Medical College, Gwalior, Madhya Pradesh, India. Phone: 0751-2626340.
E-mail: ravi_ambey97@yahoo.co.in

© 2015 International Journal of Medical Science Research and Practice available on www.ijmsrp.com